

1
2
3
4
5
6 **Glucocorticoid programming of intrauterine development**
7
8

9 **A.L. Fowden¹, O.A. Valenzuela¹, O.R. Vaughan¹, J.K. Jellyman^{1,2} & A.J. Forhead^{1,3}**
10
11

12 **¹*Centre for Trophoblast Research and Department of Physiology, Development and Neuroscience,***

13 ***University of Cambridge, Cambridge CB2 3EG, UK***

14 **²*Los Angeles Biomedical Research Institute at Harbor-University of California Los Angeles***

15 ***Medical Center, Torrance, CA 90502, USA***

16 **³*Department of Biological and Medical Sciences, Oxford Brookes University, Oxford, OX3 0BP, UK***
17
18
19
20
21
22
23
24

25 **Address for correspondence:** Abigail L. Fowden

26 Department of Physiology, Development and Neuroscience,

27 University of Cambridge,

28 Downing Street

29 Cambridge,

30 CB2 3EG, UK

31 Tel: 44 (0)1223 333855

32 Fax: 44 (0)1223 333840

33 E-mail: alf1000@cam.ac.uk
34
35

ABSTRACT

Glucocorticoids are important environmental and maturational signals during intrauterine development. Towards term, the maturational rise in fetal glucocorticoid concentrations decreases fetal growth and induces differentiation of key tissues essential for neonatal survival. When cortisol levels rise earlier in gestation as a result of suboptimal conditions for fetal growth, the switch from tissue accretion to differentiation is initiated prematurely, which alters the phenotype that develops from the genotype inherited at conception. While this improves the chances of survival should delivery occur, it also has functional consequences for the offspring long after birth. Glucocorticoids are, therefore, also programming signals that permanently alter tissue structure and function during intrauterine development to optimise offspring fitness. However, if the postnatal environmental conditions differ from those signalled *in utero*, the phenotypical outcome of early life glucocorticoid overexposure may become maladaptive and lead to physiological dysfunction in the adult. This review focusses on the role of glucocorticoids in developmental programming, primarily in farm species. It examines the factors influencing glucocorticoid bioavailability *in utero* and the effects that glucocorticoids have on the development of fetal tissues and organ systems, both at term and earlier in gestation. It also discusses the windows of susceptibility to glucocorticoid overexposure in early life together with the molecular mechanisms and long term consequences of glucocorticoid programming with particular emphasis on the cardiovascular, metabolic and endocrine phenotype of the offspring.

1. Introduction

Glucocorticoids are important stress hormones in adult animals but have a wider range of functions in the fetus. In late gestation, they act as maturational signals that ensure the fetus is mature enough to survive the transition to extra-uterine life at delivery [1]. Earlier in gestation, glucocorticoids can act as signals of suboptimal environmental conditions and modify fetal development in relation to the available resource for intrauterine growth. While improving the likelihood of survival both *in utero* and at birth, this early overexposure to glucocorticoids adapts the phenotype that develops from the genotype inherited at conception with life-long functional consequences [2-11]. Glucocorticoids are, therefore, also programming signals that permanently alter tissue structure and function during intrauterine development to optimise offspring fitness [12, 13]. Previous reviews of glucocorticoid programming have tended to concentrate on the human implications and/or the experimental studies of short lived, laboratory species like mice, rats and

guinea pigs [2-9]. In contrast, this review examines the role of glucocorticoids in developmental programming with particular emphasis on the longer-lived farm species like sheep, pigs, cattle and horses.

2. Fetal glucocorticoid exposure

There are a number of different mechanisms by which glucocorticoid concentrations can rise in the fetal circulation (Figure 1). For most of gestation, the primary source of cortisol in fetal ovine plasma is the mother [19]. Glucocorticoids cross the placenta readily by diffusion down a materno-fetal concentration gradient which exists in normal conditions in all species studied to date including pigs, cattle, sheep, pigs and horses [1,3]. Consequently, increases in maternal glucocorticoid concentrations induced by stressful conditions, such as isolation, transport, undernutrition and housing conditions, can lead to raised concentrations in the fetus [20]. However, the degree of fetal overexposure to the higher maternal glucocorticoid concentrations is minimised by the presence in the placenta of the enzyme, 11 β -hydroxysteroid dehydrogenase type 2 (11 β HSD2, Figure 1). This isoform of the enzyme converts active glucocorticoids into their inactive keto-metabolites and, hence, acts as a barrier to transplacental glucocorticoid transfer [14]. Amongst species, placental 11 β HSD2 activity appears to be positively related to the magnitude of the materno-fetal glucocorticoid gradient and is influenced by gestational age and a range of environmental factors including glucocorticoid concentrations on both sides of the placenta (Figure 1). In addition, in the hemochorial type of placenta, there are multidrug resistance transporters, which transfer xenobiotics like synthetic glucocorticoids from the trophoblast cells back into the maternal circulation (Figure 1). Abundance of these transporters are also regulated developmentally and by glucocorticoids but whether they are present in the epitheliochorial placenta of ruminants, pigs and horses remains unknown [9]. Fetal glucocorticoid concentrations can, therefore, be altered independently of maternal levels by manipulating the effectiveness of the placental barrier to materno-fetal glucocorticoid transfer. Glucocorticoid programming in early to mid-gestation, therefore, depends on the level of stress experienced by the mother during pregnancy, her HPA responses and ensuing cortisol concentration, and on the glucocorticoid permeability of the placenta.

Later in gestation when the fetal hypothalamic-pituitary-adrenal (HPA) axis has developed functionally, fetal glucocorticoid concentrations can also rise independently of maternal levels by

direct cortisol secretion from the fetal adrenal glands (Figure 1). This occurs via activation of the HPA axis in response to adverse intrauterine conditions like hypoxia and hypoglycaemia caused, for example, by cord occlusion, placental insufficiency, poor uterine perfusion or maternal alterations in dietary composition or calorie intake [20]. Development of fetal HPA responsiveness to adverse stimuli varies in timing between species and with both early glucocorticoid overexposure and repeated insults during late gestation [13,21]. Closer still to term, fetal cortisol concentrations rise naturally in the absence of adverse stimuli as part of the normal sequence of prepartum maturational events that ensure viability at birth [1; Figure 2]. The magnitude and timing of this normal prepartum cortisol surge also varies widely between species (Figure 2) and can be activated earlier than normal by poor nutritional conditions either around the time of conception or during late gestation [1,18,22,24]. Its timing is also influenced by the number of fetuses in sheep [22]. In some species like the horse, the main perinatal rise in cortisol concentrations occurs immediately after not before birth [21, 25]. The window of susceptibility to glucocorticoid programming in late gestation will, therefore, vary with species in relation to environmental conditions *in utero* and the development and responsiveness of the fetal HPA axis.

Fetal glucocorticoid overexposure can also occur as a result of clinical use of synthetic glucocorticoids like dexamethasone and betamethasone during pregnancy. These drugs are 20 times more potent than the natural hormones and are cleared more slowly from the circulation [26]. They are often used to treat conditions with an inflammatory component such as joint and respiratory problems, allergic reactions and endotoxic shock in several species [26-28]. They are also given routinely to healthy pregnant women threatened with pre-term delivery to improve neonatal viability of their infants [26]. Since the onset of labour is co-ordinated with maturation through the prepartum cortisol rise in many ruminants, synthetic glucocorticoids are also used to induce delivery of viable offspring at or near term in cattle and sheep [29-31]. Experimentally, these drugs have been used extensively in the longer-lived farm species to investigate the likely long-term physiological consequences for the human infant of antenatal glucocorticoid treatment [10,11].

The developmental effects of the glucocorticoids are determined ultimately by their bioavailability within the tissues. In turn, this depends on expression of the glucocorticoid (GR) and mineralocorticoid receptors (MR) to which the glucocorticoids bind [14]. These receptors vary in abundance between fetal tissues and with gestational age [32-34]. Their expression can also be influenced by glucocorticoid concentrations *per se* [35]. In addition, fetal tissues express 11 β HSD, both the type 2 isoform found in the placenta and the type 1 isoform which reactivates the

biologically inactive metabolites [14]. Activity of the two isoforms varies between tissues and both isoforms are regulated developmentally and by fetal oxygen, nutrient and glucocorticoid availability in a tissue specific manner [32-35]. Consequently, overexposure to glucocorticoids is determined not only systemically by the circulating concentrations but also locally within the tissues themselves. Since synthetic glucocorticoids are poorly inactivated by 11 β HSD2 and bind only to the GR [14], their bioavailability and actions can differ from those of the natural glucocorticoids.

3. Glucocorticoid programming

3.1 Glucocorticoids and fetal development

Maturationally, glucocorticoids affect development of a wide range of fetal tissues, particularly those essential for immediate neonatal survival like the lungs, liver, gut and kidneys [1,13]. During late gestation, experimental manipulation of fetal cortisol concentration in fetal sheep by fetal adrenalectomy and exogenous cortisol infusion have shown that cortisol induces changes in tissue expression of cytostructural proteins, receptors, enzymes, ion channels and growth factors [1,3]. These changes lead to alterations in tissue morphology, biochemical composition, metabolism and hormone sensitivity with functional consequences for multiple organ systems in the fetus. For example, in addition to their well established role in pulmonary maturation [1,26], glucocorticoids increase fetal blood pressure during late gestation via effects on the fetal heart and blood vessels [36]. Similarly, the fetal liver develops the capacity of gluconeogenesis close to term as a result of cortisol-induced increases in glycogen deposition and gluconeogenic enzyme activities [37]. Glucocorticoids, therefore, activate many of the physiological processes that have little or no function *in utero* but which are vital at birth like pulmonary gas exchange, hepatic gluconeogenesis and thermogenesis [1]. They also affect development of many other tissues like the brain and skeletal muscle which are important for offspring viability and fitness in the longer term [38, 39]. Consequently, delivery before adequate intrauterine exposure to rising cortisol concentrations leads to functional immaturity at birth, poor neonatal viability and/or a failure to thrive postnatally [40]. This scenario is likely to be particularly important in twin-bearing and polytocus species like sheep and pigs in which the timing of fetal HPA activation can differ between littermates.

The maturational effects of the glucocorticoids are mediated, in part, by changes in the circulating concentrations and tissue bioavailability of a range of other hormones [41]. In fetal sheep, the functioning of several endocrine systems including the HPA axis itself is affected by the prepartum

cortisol surge via changes in endocrine cell populations, enzyme activities and hormone receptor abundance (Figure 3). This leads to prepartum increases in fetal plasma concentrations of several hormones in addition to cortisol, including tri-iodothyronine (T_3), leptin and adrenaline [20]. In turn, these hormones have independent effects on development of a range of fetal tissues [41]. For instance, terminal differentiation of mononucleated cardiac myocytes to their binucleated form is initiated by the prepartum cortisol surge but depends on activation of specific tissue deiodinases and the concomitant increase in fetal T_3 bioavailability [42,45]. The changes in the set point and sensitivity of the endocrine axes induced by the prepartum cortisol surge also prepare the fetus for the new homeostatic challenges of extrauterine life. For example, the cortisol induced increases in the adrenal activity of phenylethanolamin-N-methyl transferase (PNMT) and the hepatic abundance of β -adrenoreceptors mean that neonates can secrete adrenaline in response to stressful conditions like hypoglycaemia and respond to the circulating adrenaline and produce glucose endogenously [37,46,47].

Early increases in the fetal glucocorticoid concentration also trigger tissue differentiation in the fetus [1,13]. However, in sheep, the effects of preterm cortisol administration do not entirely recapitulate the maturational changes in tissue differentiation induced by the increase in cortisol concentrations towards term. For example, adrenal PNMT abundance is increased by the prepartum cortisol surge but is decreased by cortisol administration at 100 days of gestation [46,48]. Similarly, cortisol depresses hepatic IGF-II expression at term but not earlier in gestation [49]. The transcriptome observed in the fetal lung and heart after early cortisol infusion also differs from that seen at term [50,51]. This probably reflects, in part, the ontogenic changes in tissue abundance of GR and/or other hormone receptors.

By simulating tissue differentiation, cortisol reduces tissue accretion *in utero* [1,13]. As a result, the overall rate of fetal growth declines as cortisol concentrations rise in fetal sheep towards term and in response to adverse intrauterine conditions [13]. The prepartum decline in growth rate can be prevented by fetal adrenalectomy and can be stimulated prematurely by infusing cortisol into either the fetus or mother earlier in gestation [13,52]. In several species including laboratory and farm animals, maternal administration of synthetic glucocorticoids in late gestation has also been shown to reduce offspring size, both shortly after administration and at delivery longer after the period of treatment [26,53,54; Table 1]. Similar reductions in fetal growth have been seen with administration of synthetic glucocorticoids directly to fetal sheep although the effects appear to be less pronounced than with maternal administration [53]. This suggests that the growth inhibitory

effects of synthetic glucocorticoids may be mediated, in part, by maternal metabolic changes or actions on the placenta [10, 103].

3.2 Glucocorticoids and placental development

Reductions in placental weight are seen in response to administration of both natural and synthetic glucocorticoids during mid to late gestation in sheep and other species [104]. These changes are associated with reduced expression of anti-apoptotic markers and increased expression of pro-apoptotic factors in the ovine placenta [105]. The growth inhibitory effects are more pronounced with maternal than fetal administration and tend to persist after cessation of treatment [105]. They may also be sex-linked [10]. In sheep, both maternal and fetal cortisol administration alter the gross placental morphology with proportionately fewer of the more everted placentomes [106,107]. Although the functional significance of this shift in placentome distribution remains unclear [106,108,109] placentas with fewer everted C and D type placentomes transport more glucose on a weight specific basis when fetal cortisol concentrations are high [106]. In general, glucocorticoids reduce placental glucose transport via effects on the transplacental glucose concentration gradient, placental glucose consumption and/or placental expression of the glucose transporters, dependent on the species [104]. They also reduce the active transport of amino acids across the placenta and alter placental amino acid metabolism in some species [104]. Glucocorticoid-induced changes in placental transport phenotype also vary with time both during the period of treatment and after it has ended [104]. In addition, there are alterations in the endocrine function of the placenta in response to raising glucocorticoid concentrations, which involve a wide range of hormones and changes in both their synthesis and metabolism (Figure 2). Again, these effects can be sex-linked and are often dependent on gestational age at the time of glucocorticoid exposure. For example, placental 11 β HSD2 gene expression is increased by dexamethasone administration to ewes at 30% of gestation in males alone but decreased by treatment later in pregnancy in both sexes [10,105,109].

The changes in placental endocrine and transport function induced by the prepartum cortisol surge are part of the normal sequence of events leading to labour and delivery of viable neonates [1,40]. However, their induction by glucocorticoid overexposure earlier in gestation may alter fetal growth and development independently of any direct effects of the glucocorticoids on the fetal tissues *per se* [1,3,13]. For example, the reduction in placental lactogen production and its maternal concentration in response to early glucocorticoid overexposure may alter maternal metabolism and, hence, nutrient allocation to the gravid uterus [109]. Similarly, glucocorticoid induced changes in

the production of progesterone and other progestagens may influence maternal insulin resistance and appetite with indirect consequences for intrauterine growth [110]. In addition, changes in placental phenotype induced by early glucocorticoids overexposure may persist or appear only after restoration of normal concentrations to affect fetal development long after original insult [104].

3.3 Postnatal outcomes of early overexposure to glucocorticoids

Glucocorticoid exposure *in utero* has been shown to affect organ growth and the functioning of physiological systems in the offspring after birth in sheep, pigs and cattle (Table 1). In particular, there are abnormalities in postnatal cardiovascular and metabolic function that are associated with overt hypertension and glucose intolerance by adulthood. These changes involve a wide range of tissues including the brain, blood vessels, kidneys, heart and skeletal muscle as well as several endocrine systems (Table 1). Similar findings have been made in adult rodents and humans overexposed to glucocorticoids prenatally [2-9]. Because of the central role of glucocorticoids in regulating adult cardiovascular and metabolic function, many of these studies have concentrated on programming of the HPA axis *per se* (Table 1). In sheep, pigs and cattle, early prenatal overexposure to either natural or synthetic glucocorticoids can alter both basal and stimulated cortisol concentrations postnatally and, hence, the physiological responses to homeostatic challenges (Table 1). Indeed, amongst species, maternal glucocorticoid administration during late pregnancy has been shown to programme postnatal HPA function at every level of the axis from the hippocampus to glucocorticoid bioavailability in the peripheral tissues [11]. Taken together, these studies have shown that glucocorticoids overexposure *in utero* affects the same range of tissues and cellular processes in the adult as seen in the fetus [1,3,13]. However, the specific postnatal outcomes of intrauterine glucocorticoid overexposure depend on gestational age at its onset, its severity and duration and on whether exposure was to natural or synthetic glucocorticoids (Table 1).

Natural and synthetic glucocorticoids have different programming effects. Maternal administration of cortisol but not dexamethasone at 20% of ovine pregnancy causes fasting hyperglycaemia while dexamethasone but not cortisol increases their initial insulin response to glucose administration in the adult male offspring [53]. In contrast, both cortisol and dexamethasone administration at this stage of pregnancy give rise to hypertension in the adult offspring [61]. However the mechanism by which hypertension is induced differs with cortisol increasing peripheral resistance but not cardiac output while dexamethasone enhances cardiac output but not peripheral resistance in the adult

sheep [4]. Different synthetic glucocorticoids also appear to have different programming effects (Table 1), although few, if any, studies have specifically compared the postnatal consequences of dexamethasone and betamethasone treatment *in utero*.

In sheep, glucocorticoids have been shown to have programming effects on postnatal phenotype with administration from as early as 27 days of pregnancy right up until term (Table 1). However, the specific outcomes depend on gestational age at onset of treatment (Table 1). For instance, maternal administration of a single course of dexamethasone leads to hypertension in the adult offspring when given at 27 and 80 days but not at 64 days of pregnancy [63,64]. In contrast, dexamethasone has little effect on glucose tolerance or insulin sensitivity of adult female offspring with administration at either 27 days or 64 days of gestation [64]. Similarly, maternal dexamethasone treatment early in pregnancy appears to have little effect on HPA function but, later in gestation, it decreases HPA responsiveness of the adult offspring [69,111]. Furthermore, multiple doses of betamethasone over a 14-d period in late pregnancy have subtly different effects on HPA function and glucose-insulin dynamics of the adult offspring than single doses given at the same gestational age as that at the start of the more prolonged treatment [86,87]. Longer periods of dexamethasone treatment at lower doses also have little effect relative to a single treatment at the higher, clinically relevant doses during the same period of gestation [4,112]. Consequently, the dose of synthetic glucocorticoid administered as well as its duration and timing in pregnancy is important in determining the phenotypical outcome.

Another factor influencing the apparent extent of programming is the postnatal age at which the outcomes are assessed (Table 1). Some glucocorticoid-programmed changes in postnatal growth and physiological function are apparent immediately after birth while others only become evident later in life as the animal ages, reaches key life course events like weaning, puberty or pregnancy or experiences adverse conditions after birth [113; Table 1]. For example, hypertension is not seen in the neonatal lamb after intrauterine dexamethasone overexposure, although there are changes in the baroreflexes indicative of resetting of the neural mechanisms of blood pressure control even at this early stage of postnatal life [56]. Hypertension is evident at 4 months of age at about the time weaning is complete and becomes progressively more pronounced with increasing age thereafter [57]. Overall, the experimental studies suggest that glucocorticoid-programmed metabolic dysfunction appears later in ovine life than the cardiovascular abnormalities and is often not detected until adulthood (Table 1). Metabolic changes may also only be detected in one sex (Table

1). For instance, altered glucose-insulin dynamics are seen in 4-5 year-old male but not female offspring overexposed to dexamethasone at 27 days of gestation [55,64]. In addition, there is emerging evidence in rodents and other species that maternal diet and pre-existing conditions such as intrauterine growth restriction can influence the fetoplacental responses to glucocorticoid administration, which, in turn, are likely to affect programming of postnatal phenotype [114,115].

3.4 Developmental windows of glucocorticoid programming

Collectively, the experimental studies summarised in Table 1 suggest that there are specific stages in development when glucocorticoid overexposure is most likely to result in an altered postnatal phenotype. The first window of susceptibility is probably during pre-implantation development when lineage specification occurs and cells are segregated into trophectoderm and inner cell mass. Certainly, undernutrition during this period of pregnancy has effects on development of the fetal HPA axis and other organ systems much later in gestation [18,24]. The second vulnerable period for glucocorticoid programming is during organogenesis which occurs between days 7 and 30 of gestation in sheep embryos. This also covers the period of implantation and formation of the ovine placenta [10]. Indeed, compromised development of the metanephric kidney is likely to be a significant contributory factor in the hypertension seen in adult offspring of ewes treated with dexamethasone at 27 days of gestation [4]. After completing organogenesis, there is a relatively long period of gestation when the fetus is gaining mass and developing the neural and endocrine mechanisms regulating homeostasis. During this period, excess glucocorticoids appear to act by changing the kinetics of the cell cycle to slow growth and set the responsiveness of the regulatory mechanisms. When tissues have developed sufficient GR or at critical concentrations or duration of exposure, glucocorticoids can switch the cell cycle from proliferation to differentiation prematurely, with permanent effects on total cell number and/or the balance of different cell types within an organ [13]. For example, cortisol induced differentiation of cardiomyocytes from the mononucleated form, which can divide, to the binucleated type, which cannot, means that cell number is fixed and that cardiac growth depends primarily on cell hypertrophy rather than hyperplasia thereafter [42]. Finally, in some species, there appears to be a window of susceptibility to glucocorticoid programming in the period immediately after birth, which may be particularly important in species like the horse in which terminal differentiation of tissues is not complete at birth [21]. Neonatally, cortisol overexposure may occur either directly due to sickness or maladaptation *ex utero* or indirectly via changes in milk composition and its glucocorticoid content as a result of maternal stress or abnormal mammary development [116,117]. Certainly,

experimental overexposure of healthy newborn foals to cortisol for 5 days after birth alters both HPA and pancreatic β cell function later in life (Table 1).

3.5 Molecular mechanisms of glucocorticoid programming

At the molecular level, there appears to be two broad mechanisms by which glucocorticoids act to programme development. First, bound to their receptors, they may act as enhancer binding proteins that activate or repress expression of genes via interaction with glucocorticoid response elements in the promotor or other regulatory regions of the genome [118]. With genes that trigger key developmental stages, their altered expression at inappropriate times in the normal sequence of events may have permanent effects on the subsequent pattern of development. This type of glucocorticoid-induced change in expression of specific genes may occur either early in development, for example during cell lineage specification and mitochondrial biogenesis, or later in gestation during differentiation of sub-populations of cells within tissues such as the liver or endocrine pancreas. The outcomes of these discrete gene expression events may, therefore, be global in terms of cell metabolism and oxidative stress or specific to certain tissues or cell types. In rodents, increased oxidative stress is a common feature of glucocorticoid programming along with changes to the relative numbers of the different endocrine cell types within the Islets of Langerhans [2-9].

Secondly, glucocorticoids may alter the epigenome with more long term consequences for gene expression throughout life [119]. Changes in DNA methylation and histone modifications have been observed both globally and in specific tissues in postnatal offspring glucocorticoid overexposed *in utero* [35,119]. In particular, there are tissue specific changes in the methylation status of the regulatory regions of the GR gene which influence expression of these receptors and, hence, postnatal glucocorticoid responsiveness [33, 118]. Maternal dexamethasone administration during pregnancy has been shown to reduce placental transport of folate required for one carbon metabolism and DNA methylation while, conversely, dietary supplementation with folate ameliorates, in part, the feto-placental growth restriction induced by this treatment [115,121]. In addition, there is emerging evidence for postnatal changes in expression of various non-coding and microRNAs after early life glucocorticoid overexposure [39]. Glucocorticoids, therefore, affects the developing epigenome through a number of different routes with dynamic consequences for epigenetic marks throughout the lifespan of the animal. However, to date, most of the information

about the molecular mechanisms of glucocorticoid programming has been derived from studies in rodents and guinea pigs so the extent to which they apply to farm species remains unclear.

4. Conclusions

Glucocorticoids have a number of roles during intrauterine and early neonatal development. Not only are they essential for normal maturation close to term, they also act as important signals of environmental compromise earlier in gestation. The glucocorticoid triggered switch from tissue accretion to differentiation improves offspring fitness by maximising the chances of the fetus surviving into adulthood. Prenatally, early activation of this switch ensures that fetal growth is commensurate with the nutrient supply *in utero* and that fetal tissues are sufficiently mature to function *ex utero* should delivery occur. Postnatally, the glucocorticoid-induced adaptations in phenotype and, particularly the resetting of the homeostatic control mechanisms, will help the offspring to thrive in a postnatal environment matching that signalled to it *in utero*. However, when pre- and post-natal environments are mismatched in laboratory species, the glucocorticoid-induced changes in offspring phenotype can become maladaptive and lead to early onset of cardiometabolic dysfunction characteristic of old age [2-9]. Recent findings have also shown that the effects of early life glucocorticoid overexposure can persist inter-generationally with changes in F2 placental phenotype and HPA function after dexamethasone treatment of their pregnant grandmothers [122-125]. This raises the possibility that glucocorticoids may also have an important evolutionary role in the transgenerational inheritance of phenotypical traits. However, the extent to glucocorticoid overexposure during early development influences lifespan and transgenerational inheritance in longer lived farm species remains largely unknown.

5. References

- [1] Fowden AL, Li J, Forhead AJ. Glucocorticoids and the preparation for life after birth: are there long term consequences of the life insurance? *Proc Nutr Soc* 1998;57:113-122.
- [2] Seckl JR. Prenatal glucocorticoids and long term programming. *Eur J Endocrinol* 2004;Suppl 3 U49-U62.
- [3] Fowden AL, Forhead AJ. Endocrine mechanisms of intrauterine programming. *Reproduction* 2004;127:515-526.
- [4] Moritz KM, Boon WM, Wintour EM. Glucocorticoid programming of adult disease. *Cell Tissue Res*. 2005;322:81-88.
- [5] Khulan B, Drake AJ. Glucocorticoids as mediators of developmental programming effects. *Best Prac Res Clin Endocrinol Metab* 2012;26:689-700.
- [6] Painter RC, Roseboom TJ, de Rooij SR. Long-term effects of prenatal stress and glucocorticoid exposure. *Birth Defects Res* 2012;96:315-324.
- [7] Reynolds RM. Glucocorticoid excess and the developmental origins of adult disease: two decades of testing the hypothesis. *Psychoneuroendocrinology* 2013;38:1-11.
- [8] McKinley CJD, Dalziel SR, Harding JE. Antenatal glucocorticoids: where are we after forty years? *J Dev Orig Health Dis*. 2015;6:127-142
- [9] Moisiadis VG, Matthews SG. Glucocorticoids and fetal programming Part 1: Outcomes. *Nat Rev Endocrinol*. 2014;10:391-402.
- [10] Braun T, Challis JR, Newnham JP, Sloboda DM. Early-life glucocorticoid exposure: the hypothalamic-pituitary-adrenal axis, placental function and long-term disease risk. *Endocr Rev*. 2013;34:885-916.
- [11] Jellyman JK, Valenzuela OA, Fowden AL. Glucocorticoid programming of the hypothalamic-pituitary-adrenal axis and metabolic function: Animal studies from mouse to horse. *J Anim Sci*. 2015;93:3245-3260.
- [12] Fowden AL, Moore T. Maternal-fetal resource allocation: co-operation and conflict. *Placenta* 2012;33 Suppl. 2:e11-e15.
- [13] Fowden AL, Forhead AJ. Glucocorticoids as regulatory signals during intrauterine development. *Exp Physiol*. DOI:10.1113/EP085212.
- [14] Chapman K, Holmes M, Seckl JR. 11 β -hydroxysteroid dehydrogenase: intracellular gate-keepers of tissue glucocorticoid action. *Physiol Rev*. 2013;93:1139-1206.
- [15] Braun T, Li S, Moss TJ, Connor KL, Doherty DA, Nitsos I, Newnham JP, Challis JR, Sloboda DM. Differential appearance of placentomes and expression of prostaglandin H synthase type 2 placentome subtypes after betamethasone treatment of sheep late in gestation. *Placenta* 2011;32:295-303.

446 [16] Kerzner LS, Stonestreet BS, Wu KY, Sadowska G, Malee, MP. Antenatal dexamethasone: effects
447 on ovine placental 11 beta-hydroxysteroid dehydrogenase type 2 expression and fetal growth.
448 *Pediatr Res.* 2002;52:706-712.

449 [17] Clarke K, Ward JW, Forhead AJ, Giussani DA, Fowden AL. Regulation of 11 β -hydroxysteroid
450 dehydrogenase type 2 (11 β -HSD2) activity in ovine placenta by fetal cortisol. *J Endocrinol.*
451 2002;172:527-534.

452 [18] McMullen S, Ogersby JC, Thurston LM, Gadd TS, Wood PJ, Wathes DC, Michael AE. Alterations in
453 11 beta-hydroxysteroid dehydrogenase (11 β HSD) activities and fetal cortisol:cortisone ratios
454 induced by nutritional restriction prior to conception and at defined stages of gestation in ewes.
455 *Reproduction* 2004;127:717-725.

456 [19] Hennessy DP, Coghlan JP, Hardy KJ, Scoggins BM, Wintour EM. The origin of cortisol in the blood
457 of fetal sheep. *J Endocrinol.* 1982;95:71-79.

458 [20] Fowden AL, Forhead AJ. Hormones as epigenetic signals in developmental programming. *Exp*
459 *Physiol.* 2009;94:607-625.

460 [21] Rouffle J, Dalle M, Touraire C, Barlet J-P, Delost P. Sodium intake by pregnant cows and plasma
461 aldosterone and cortisol concentrations in the fetus during late pregnancy. *J Dairy Sci* 1990;73:1762-
462 1765.

463 [22] Vannucchi CI, Rodrigues JA, Silva LCG, Lucio CF, Veiga GAL, Furtado PV, Oliveira CA, Nichi M.
464 Association between birth conditions and glucose and cortisol profiles of periparturient dairy cows
465 and neonatal calves. *Vet Rec* 2012;176:358

466 [23] Fowden AL, Silver M. Comparative development of the pituitary-adrenal axis in the fetal foal
467 and lamb. *Reprod Domest Animal* 1995;30:170-177.

468 [24] Edwards LJ, McMillen IC. Impact of maternal undernutrition during the periconceptional period,
469 fetal number, and fetal sex on the development of the hypothalamic-pituitary adrenal axis in sheep
470 during late gestation. *Biol Reprod.* 2002;65:1562-1569.

471 [25] Silver M, Cash RSG, Dudan F, Fowden AL, Knox J, Ousey JC, Rosedale PD. Postnatal
472 adrenocortical activity in relation to plasma ACTH and catecholamine levels in term and premature
473 foals. *Equine Vet J.* 1984;16:278-286.

474 [26] Forhead AJ, Fowden AL. The consequences for preterm infants of antenatal glucocorticoid
475 treatment. 2011 In *Birth rites and rights* Ed. F. Ebtehaj, J. Herring, M.H. Johnson & M. Richards. pp
476 29-149. Hart Publishing.

477 [27] Leclere M, Lefebvre-Lavoie J, Beauchamp G, Lavoie JP. Efficacy of prednisone and
478 dexamethasone in horses with recurrent airway obstruction in the presence of continuous antigen
479 exposure. *Equine Vet J.* 2010;42:316-321.

480 [28] Johnson PJ, Slight SH, Ganjam VK, Kreeger JM. Glucocorticoids and laminitis in the horse. *Vet*
481 *Clin North Am Equine Pract.* 2002;18:219-236.

482 [29] Kastelic JP, Cook RB, McMahon LR, McAllister TA, McClelland LA, Cheng KJ. Induction of
483 parturition in ewes with dexamethasone or dexamethasone and cloprostenol. *Can Vet J.*
484 1996;37:101-102.

485 [30] Mansell PD, Cameron AR, Taylor DP, Malmo J. Induction of parturition in dairy cattle and its
486 effects on health and subsequent lactation and reproductive performance. *Aust Vet J.* 2006;84:312-
487 316.

488 [31] Shenavai S, Preissing S, Hoffman B, Dilly M, Pfarrer C, Ozalp GR, Caliskan C, Seyrek-Intas K,
489 Schular G. Investigations into the mechanisms controlling parturition in cattle. *Reproduction*
490 2012;144:279-292.

491 [32] Shang H, Meng W, Sloboda DM, Li S, Ehlich L, Plagemann A, Dudenhausen JW, Henrich W,
492 Newnham JP, Challis JR, Braun T. Effects of maternal dexamethasone treatment in pregnancy on
493 glucocorticoid receptors in the ovine placenta. *Reprod Sci.* 2015;22:535-544.

494 [33] Sloboda DM, Moss TJ, Li S, Matthews SG, Challis JR, Newnham JP. Expression of glucocorticoid
495 receptor, mineralocorticoid receptor, and 11 beta-hydroxysteroid dehydrogenase 1 and 2 in the fetal
496 and postnatal ovine hippocampus: ontogeny and effects of prenatal glucocorticoid exposure. *J*
497 *Endocrinol.* 2008;197:213-220.

498 [34] Brown RW, Diaz R, Robson AC, Kotelevtsev YV, Mullins JJ, Kaufman MH, Seckl JR . The ontogeny
499 of 11 β -hydroxysteroid dehydrogenase type 2 and mineralocorticoid receptor gene expression reveal
500 intricate control of glucocorticoid action in development. *Endocrinol* 1996;137:794-797.

501 [35] Moisiadis VG, Matthews SG. Glucocorticoids and fetal programming Part 2: Mechanisms. *Nat*
502 *Rev Endocrinol.* 2014;10:403-411.

503 [36] Unno N, Wong GH, Jenkins SL, Wentworth RA, Ding XY, Li C, Robertson SS, Smotherman WP,
504 Nathanielsz PW. Blood pressure and heart rate in the ovine fetus: ontogenic changes and effects of
505 fetal adrenalectomy. *Am J Physiol.* 1999;276:H248-H256.

506 [37] Fowden AL, Mundy L, Silver M. Developmental regulation of glucogenesis in the sheep fetus
507 during late gestation. *J Physiol.* 1998;508:937-947.

508 [38] Peffer M, Zhang JY, Umfrey L, Rudine A, Monaghan P, Defranco DB. Minireview: The impact of
509 antenatal therapeutic synthetic glucocorticoids on the developing brain. *Mol Endocrinol.*
510 2015;29:658-666.

511 [39] Bouret S, Levin BE Ozanne SE. Gene-environment interactions controlling energy and glucose
512 homeostasis and the developmental origin of obesity. *Physiol Rev.* 2015;95:47-82.

513 [40] Silver M. Prenatal maturation, the timing of birth and how it may be regulated in domestic
514 animals. *Quart J Exp Physiol* 1990;75:285-307.

515 [41] Fowden AL, Forhead AJ. Endocrine interactions in the control of fetal growth. *In* *Maternal and*
516 *Child Nutrition.* 2013 Ed. Bhatia, J., Bhutta, Z. & Kahlan, S.C. Nestlé Nutrition Workshop Series 74 91-
517 102.

518 [42] Thornburg K, Jonker S, O'Tierney P, Chattergoon N, Louey S, Faber J, Giraud G. Regulation of the
519 cardiomyocyte population in the developing heart. *Prog Biophy Mol Biol.* 2011;106:289-299.

520 [43] Forhead AJ, Jellyman JK, De Blasio MJ, Johnson E, Giussani DA, Broughton Pipkin F, Fowden AL.
521 Maternal dexamethasone treatment alters tissue and circulating components of the renin-
522 angiotensin system in the pregnant ewe and fetus. *Endocrinol.* 2015;156:3038-3046.

523

524 [44] De Blasio MJ, Boije M, Vaughan, OR, Bersteins, BS, Davies, KL, Plein, A, Kempster SL, Smith GCS,
525 Charnock-Jones DS, Blache D, Wooding FBP, Giussani DA, Fowden AL, Forhead AJ. Developmental
526 expression and glucocorticoid control of leptin receptors in the fetal ovine lung. PLOS ONE
527 2015;10:e136115.

528 [45] Forhead AJ, Fowden AL. Thyroid hormones in fetal growth and *prepartum* maturation. J
529 Endocrinol. 2014;221:R87-103.

530

531 [46] Comline RS, Silver M. Activity of the adrenal medulla before and after birth. Brit Med Bull.
532 1966;22:16-20.

533 [47] Fowden AL, Apatu, RSK, Silver M. The glucogenic capacity of the fetal pig: developmental
534 regulation by cortisol. Exp Physiol. 1995;80:457-467.

535 [48] Adams MB, Ross JT, Butler TG, McMillen IC. Glucocorticoids decrease phenylethanolamine N-
536 methyltransferase mRNA expression in the immature foetal sheep adrenal. J Neuroendocrinol.
537 1999;11:569-575.

538 [49] Li J, Saunders JC, Gilmour RS, Silver M, Fowden AL. Insulin-like growth factor II mRNA expression
539 in fetal tissues of the sheep during late gestation: effects of cortisol. *Endocrinol.* 1993;132:2083-
540 2089.

541 [50] McGillick EV, Orgeig S, McMillen IC, Morrison JL. The fetal sheep lung does not respond to
542 cortisol infusion during the late canalicular phase of development. Physiol Resp. 2013;1:e00130.

543 [51] Richards EM, Wood CE, Rabaglino MB, Antolic A, Keller-Wood M. Mechanisms for adverse
544 effects of late gestational increases in maternal cortisol on the heart revealed by transcriptomic
545 analyses of the fetal septum. Physiol Genomics. 2014;46:547-559.

546 [52] Fowden AL, Szemere J, Hughes P, Gilmour RS, Forhead A.J. The effects of cortisol on the growth
547 rate of the sheep fetus during late gestation. J Endocrinol. 1996;151:97-105.

548 [53] Newnham JP, Evans SF, Godfrey M, Huang W, Ikegami M, Jobe A. Maternal, but not fetal,
549 administration of corticosterone restricts fetal growth. J Matern Fetal Med. 1999;8:81-87.

550 [54] Kutzler MA, Ruane EK, Coksaygan, Vincent SE, Nathanielsz PW. Effects of three courses of
551 maternally administered dexamethasone at 0.7, 0.75 and 0.8 of gestation on prenatal and postnatal
552 growth in sheep. Pediatrics 2004;113:313-319.

553 [55] De Blasio MJ, Dodic M, Jefferies AJ, Moritz KM, Wintour EM, Owens JA. Maternal exposure to
554 dexamethasone or cortisol in early pregnancy differentially alters insulin secretion and glucose
555 homeostasis in adult male sheep offspring. Am J Physiol Endocrinol Metab. 2007;293:E75-E82.

556 [56] Segar JL, Roghair RD, Segar EM, Bailey MC, Scholz TD, Lamb FS. Early gestation dexamethasone
557 alters baroreflex and vascular responses in newborn lambs before hypertension. Am J Physiol Regul
558 Integr Comp Physiol. 2006;291:R481-R488.

559 [57] Dodic M, Abouantoun T, O'Connor A, Wintour EM, Moritz KM. Programming effects of short
560 prenatal exposure to dexamethasone in sheep. Hypertension. 2002;40:729-734.

561 [58] Dodic M, Samuel C, Moritz K, Wintour EM, Morgan J, Grigg L, Wong J. Impaired cardiac
562 functional reserve and left ventricular hypertrophy in adult sheep after prenatal dexamethasone
563 exposure. Circ Res. 2001;89:623-629.

564 [59] Moritz KM, De Matteo R, Dodic M, Jefferies AJ, Arena D, Wintour EM, Probyn ME, Bertram JF,
565 Singh RR, Zanini S, Evans RG. Prenatal glucocorticoid exposure in the sheep alters renal development
566 in utero: implications for adult renal function and blood pressure control. *Am J Physiol Regul Integr*
567 *Comp Physiol.* 2011;301:R500-R509.

568 [60] Wintour EM, Moritz KM, Johnson K, Rocardo S, Samuel CS, Dodic M. Reduced nephron number
569 in adult sheep; hypertension as a result of prenatal glucocorticoid treatment. *J Physiol.*
570 2003;549:929-935.

571 [61] Roghair RD, Miller FJ, Scholz TD, Lamb FS, Segar JL. Endothelial superoxide production is altered
572 in sheep programmed by early gestation dexamethasone exposure. *Neonatology* 2008;93:19-27.

573 [62] Von Bergen NH, Koppenhafer SL, Spitz DR, Volk KA, Patel SS, Roghair RD, Lamb FS, Segar JL,
574 Scholz TD. Fetal programming alters reactive oxygen species production in sheep cardiac
575 mitochondria. *Clin Sci.* 2009;116:659-668.

576 [63] Dodic M, McAlinden AT, Jefferies AJ, Wintour EM, Cock ML, May CN, Evans RG, Moritz KM.
577 Differential effects of prenatal exposure to dexamethasone or cortisol on circulatory control
578 mechanisms mediated by angiotensin II in the central nervous system of adult sheep. *J Physiol.*
579 2006;571:651-660.

580 [64] Gatford KL, Wintour EM, De Balsio MJ, Owens JA, Dodic M. Differential timing for programming
581 of glucose homeostasis, sensitivity to insulin and blood pressure by in utero exposure to
582 dexamethasone in sheep. *Clin Sci.* 2000;98:553-560.

583 [65] Li S, Moss TJM, Nitsos I, Matthews SG, Challis JRG, Newnham JP, Sloboda DM. The impact of
584 maternal synthetic glucocorticoid administration in late pregnancy on fetal and early neonatal
585 hypothalamic-pituitary-adrenal axis regulatory genes is dependent upon dose and gestational age at
586 exposure. *J Dev Orig Health Dis.* 2013;4:77-89.

587 [66] Li S, Sloboda DM, Moss TJM, Nitsos I, Polglase GR, Doherty DA, Newnham JP, Challis JRG, Braun
588 T. Effects of glucocorticoid treatment given in early or late gestation on growth and development in
589 sheep. *J DOHaD* 2013;4:146-156.

590 [67] Li S, Nitsos I, Polglase GR, Braun T, Newnham JP, Challis JR. The effects of dexamethasone
591 treatment in early gestation on hypothalamic-pituitary-adrenal responses and gene expression at 7
592 months of postnatal age in sheep. *Reprod Sci.* 2012;19:260-270.

593 [68] Long NM, Shasa DR, Ford SP, Nathanielsz PW. Growth and insulin dynamics in two generations
594 of female offspring of mothers receiving a single course of synthetic glucocorticoids. *Am J Obstet*
595 *Gynecol.* 2012;207:203e1-8.

596 [69] Long NM, Ford SP, Nathanielsz PW. Multigenerational effects of fetal dexamethasone exposure
597 on the hypothalamic-pituitary-adrenal axis of first- and second-generation female offspring. *Am J*
598 *Obstet Gynecol.* 2013;208:217e1-8.

599 [70] Bispham J, Heasman L, Clarke L, Ingleton PM, Stephenson T, Symonds ME. Effect of maternal
600 dexamethasone treatment and ambient temperature on prolactin receptor abundance in brown
601 adipose and hepatic tissue in the foetus and new-born lamb. *J Neuroendocrinol.* 1999;11:849-856.

602 [71] Berry MJ, Jacuiery AL, Oliver MH, Harding JE, Bloomfield FH. Antenatal corticosteroid exposure
603 at term increases adult adiposity: an experimental study in sheep. *Acta Obstet Gynecol Scand.*
604 2013;92:862-865.

605 [72] Shaltout HA, Chappell MC, Rose JC, Diz DI. Exaggerated sympathetic mediated responses to
606 behavioural or pharmacological challenges following antenatal betamethasone exposure. *Am J*
607 *Physiol Endocrinol Metab.* 2011;300:E979-E985.

608 [73] Su Y, Carey LC, Rose JC, Pulgar VM. Antenatal glucocorticoid exposure enhances the inhibition of
609 adrenal steroidogenesis by leptin in a sex-specific fashion. *Am J Physiol Endocrinol Metab.*
610 2013;304:E1404-E1411.

611 [74] Shaltout HA, Figueroa JP, Rose JC, Diz DI, Chappell MC. Alterations in circulatory and renal
612 angiotensin-converting enzyme and angiotensin-converting enzyme 2 in fetal programmed
613 hypertension. *Hypertension* 2009;53:404-408.

614 [75] Figueroa JP, Rose JC, Massmann GA, Zhang J, Acuna G. Alterations in fetal kidney development
615 and elevations in arterial blood pressure in young adult sheep after clinical doses of antenatal
616 glucocorticoids. *Pediatr Res.* 2005;58:510-515.

617 [76] Gwathmey TM, Shaltout HA, Rose JC, Diz DI, Chappell MC. Glucocorticoid-induced fetal
618 programming alters the functional complement of angiotensin receptor subtypes within the kidney.
619 *Hypertension* 2011;57:620-626.

620 [77] Kantorowicz L, Valego NK, Tang L, Figueroa JP, Chappell MC, Carey LC, Rose JC. Plasma and renal
621 renin concentrations in adult sheep after prenatal betamethasone exposure. *Reprod Sci.*
622 2008;15:831-838.

623 [78] Tang L, Bi J, Valego N, Carey L, Figueroa J, Chappell M, Rose JC. Prenatal betamethasone
624 exposure alters renal function in immature sheep: sex differences in effects. *Am J Physiol Regul*
625 *Integr Comp Physiol.* 2010;299:R793-R803.

626 [79] Contag SA, Bi J, Chappell MC, Rose JC. Developmental effect of antenatal exposure to
627 betamethasone on renal angiotensin II activity in the young adult sheep. *Am J Physiol Renal Physiol.*
628 2010;298:F847-F856.

629 [80] Zhang J, Massmann GA, Rose JC, Figueroa JP. Differential effects of clinical doses of antenatal
630 dexamethasone on nephron endowment and glomerular filtration rate in adult sheep. *Reprod Sci.*
631 2010;17:186-195.

632 [81] Eckman DM, Kerr BA, Fuloria M, Simandle SA, Watt SE, Rose JC, Figueroa JP. Antenatal
633 betamethasone alters vascular reactivity in adult female ovine cerebral arteries. *Pediatr Res.*
634 2010;68:344-348.

635 [82] Moss TJ, Doherty DA, Nitsos I, Sloboda DM, Harding R, Newnham JP. Effects into adulthood of
636 single or repeated antenatal corticosteroid in the sheep. *Am J Obstet Gynecol.* 2005;192:146-152.

637 [83] Pedrana G, Viotti MH, Souza E, Sloboda DM, Martin GB, Cavestrany D, Ortega HH. Apoptosis-
638 related protein expression during pre- and post-natal testicular development after administration of
639 glucocorticoids *in utero* in the sheep. *Reprod Domest Anim.* 2013;48:795-802.

640 [84] Moss TJM, Sloboda DM, Gurrin LC, Harding R, Challis JRG, Newnham JP. Programming effects in
641 sheep of prenatal growth restriction and glucocorticoid exposure. *Am J Physiol Regul Integr Comp*
642 *Physiol.* 2001;281:R960-R970.

643 [85] Gatford KL, Owens JA, Li S, Moss TJ, Newnham JP, Challis JR, Sloboda DM. Repeated
644 betamethasone treatment of pregnant sheep programs persistent reductions in circulating IGF-I and
645 IGF-binding proteins in progeny. *Am J Physiol Endocrinol Metab.* 2008;295:E170-E178.

646 [86] Sloboda DM, Moss TJ, Gurrin LC, Newnham JP, Challis JR. The effect of prenatal betamethasone
647 administration on postnatal ovine hypothalamic-pituitary-adrenal function. *J Endocrinol.*
648 2002;172:71-81.

649 [87] Sloboda DM, Moss TJ, Li S, Doherty DA, Nitsos I, Challis JR, Newnham JP. Prenatal
650 betamethasone exposure results in pituitary-adrenal hyporesponsiveness in adult sheep. *Am J*
651 *Physiol Endocrinol Metab.* 2007;292:E61-E71.

652 [88] Sloboda DM, Moss TJ, Li S, Doherty DA, Nitsos I, Challis JR, Newnham JP. Hepatic glucose
653 regulation and metabolism in adult sheep: effects of prenatal betamethasone. *Am J Physiol*
654 *Endocrinol Metab.* 2005;289:E271-E278.

655 [89] Alm CC, Sullivan JJ, First NL. The effect of a corticosteroid (dexamethasone), progesterone,
656 oestrogen and prostaglandin F2alpha on gestation length in normal and ovariectomized mares. *J*
657 *Reprod Fertil.* 1975;Suppl 23:637-640.

658 [90] Ousey JC, Kolling M, Kindahl H, Allen WR. Maternal dexamethasone treatment in late gestation
659 induces precocious fetal maturation and delivery in healthy Thoroughbred mares. *Equine Vet J.*
660 2011;43:424-429.

661 [91] Corner RA, Kenyon PR, Stafford KJ, West DM, Oliver MH. The effect of different types of
662 stressors during mid- and late pregnancy on lamb weight and body size at birth. *Animal* 2010;4:2065-
663 2070.

664 [92] Roussel S, Hemsworth PH, Boissy A, Duvaux-Ponter C. Effects of repeated stress during
665 pregnancy in ewes on the behavioural and physiological responses to stressful events and birth
666 weight of their offspring. *Appl Anim Behav Sci.* 2004;85:259-276.

667 [93] Mack LA, Lay DC, Eicher SD, Johnson AK, Richert BT, Pajor EA. Growth and reproductive
668 development of male piglets are more vulnerable to midgestation maternal stress than that of
669 female piglets. *J. Anim Sci.* 2014;92:530-548.

670 [94] Brussow K-P, Schneider F, Kanitz E, Otten W, Tuchscherer M. Alteration of reproductive
671 hormone levels in pregnant sows induced by repeated ACTH application and its possible influence on
672 pre- and post-natal hormone secretion of piglets. *J Reprod Dev.* 2005;51:133-142.

673 [95] Kanitz E, Otten W, Tuchscherer M. Changes in endocrine and neurochemical profiles in neonatal
674 pigs prenatally exposed to increase maternal cortisol. *J Endocrinol.* 2006;191:207-220.

675 [96] Haussmann MF, Carroll JA, Weesner GD, Daniels MJ, Matteri RL, Lay DC. Administration of ACTH
676 to restrained pregnant sows alters their pigs' hypothalamic-pituitary-adrenal (HPA) axis. *J Anim Sci.*
677 2000;78:2399-2411.

678 [97] Jarvis SJ, Moinard C, Robson SK, Baxter E, Ormandy E, Douglas AJ, Seckl JR, Russell JA, Lawrence
679 AB. Programming of the offspring of the pig by prenatal stress: Neuroendocrine activity and
680 behaviour. *Horm Behav.* 2006;49:68-80.

681 [98] Lay DC, Randel RD, Friend TH, Jenkins OC, Neuendorff DA, Bushong DM, Lanier EK, Bjorge MK.
682 Effects of prenatal stress on suckling calves. *J Anim Sci.* 1997;75:3143-3151.

683 [99] McGowan JE, Sysyn G, Petersson KH, Sadowska GB, Mishra OM, Delivoria-Papadopoulos M,
684 Stonestreet BS. Effect of dexamethasone treatment on maturational changes in the NMDA receptor
685 in sheep brain. *J Neuroscience.* 2000;20:7424-7429.

686 [100] Jellyman JK, Allen VL, Holdstock NB, Fowden AL. Glucocorticoid over-exposure in neonatal life
687 alters pancreatic β cell function in newborn foals. J Anim Sci. 2013;91:104-110.
688

689 [101] Jellyman JK, Allen VL, Forhead AJ, Holdstock NB, Fowden AL. (2012). Hypothalamic-pituitary-
690 adrenal axis function in pony foals after neonatal glucocorticoid overexposure. Equine Vet J.
691 2012;44:Suppl 41, 38-42.
692

693 [102] Jellyman JK, Valenzuela OA, Allen VL, Forhead AJ, Holdstock NB, Fowden AL. (2015). Neonatal
694 glucocorticoid overexposure programmes pituitary-adrenal function in ponies. Dom Anim
695 Endocrinol. 2015;50:45-49.

696 [103] Vaughan OR, Forhead AJ & Fowden AL. Glucocorticoids and placental programming. In The
697 Placenta and Human Developmental Programming, 2011. Eds. Burton, G.J., Barker, D.J.P., Moffett,
698 A., & Thornburg, K.L. pp 175-186. Cambridge University Press, Cambridge.

699 [104] Fowden AL, Forhead AJ, Sferruzzi-Perri AN, Burton GJ, Vaughan OR. (2015). Endocrine
700 regulation of placental phenotype. *Placenta* 2015;36:Suppl. 1, *Trophoblast Research* 29 S50-S59.

701 [105] Braun T, Meng W, Shang H, Li S, Sloboda DM, Ehrlich L, Lange K, Henrich W, Dudenhausen JW,
702 Plagemann A, Newnham JP, Challis JR. Early dexamethasone treatment induced placental apoptosis
703 in sheep. Reprod Sci. 2015 DOI:10.1177/1933719114542028.

704 [106] Ward JW, Forhead AJ, Wooding FBP, Fowden AL (2006). Functional significance and cortisol
705 dependence of the gross morphology of ovine placentomes during late gestation. Biol Reprod.
706 2006;74:137-145.

707 [107] Jensen EC, Gallaher BW, Breier BH, Harding JE. The effect of chronic maternal cortisol infusion
708 on the late-gestation fetal sheep. J Endocrinol. 2002;174:27-36.

709 [108] Vonnahme KA, Arndt WJ, Jihson ML, Borowicz PP, Reynolds LP. Effect of morphology on
710 placentome size, vascularity and vasoreactivity in late pregnant sheep. Biol Reprod. 2008;79:976-
711 982.

712 [109] Braun T, Li S, Moss TJ, Newnham JP, Challis JR, Gluckman PD, Sloboda DM. Maternal
713 betamethasone administration reduces binucleate cell number and placental lactogen in sheep. J
714 Endocrinol. 2007;194:337-347.

715 [110] Vejrazkova D, Vcelak J, Vankova M, Lukasova P, Bradnova O, Halkova T, Kancheva R, Bendlova
716 B. Steroids and insulin resistance during pregnancy. J Steroid Biochem Mol Biol. 2014;139:122-129.

717 [111] Dodic M, Peers A, Moritz K, Hantzis V, Wintour EM. No evidence for HPA reset in adult sheep
718 with high blood pressure due to short prenatal exposure to dexamethasone. Am J Physiol Regul
719 Integr Comp Physiol. 2002;282:R343-R350.

720 [112] Moritz K, Butkus A, Hantzis V, Peers A, Wintour EM, Dodic M. Prolonged low-dose
721 dexamethasone, in early gestation, has no long term-term deleterious effect on normal ovine
722 fetuses. Endocrinol. 2002;143:1159-1165.

723 [113] Wada H. Glucocorticoids: mediators of vertebrate ontogenetic transitions. Gen Comp
724 Endocrinol. 2008;156:441-453.
725

- [114] Miller SL, Supramaniam VG, Jenkin G, Walker DW, Wallace EM. Cardiovascular responses to maternal betamethasone administration in the intrauterine growth-restricted ovine fetus. *Am J Obstet Gynecol.* 2009;201:612e1-8.
- [115] Zhou L, Zhang A, Wang K, Zhou Q, Duan T. Folate ameliorates dexamethasone-induced fetal and placental growth restriction potentially via improvement of trophoblast migration. *Int J Clin Exp Pathol.* 2015;8:3008-3014.
- [116] Henderson JJ, Hartmann PE, Moss TJ, Doherty DA, Newnham JP. Disrupted secretory activation of the mammary gland after antenatal glucocorticoid treatment in sheep. *Reproduction* 2008;136:649-655.
- [117] Hinde K, Skibel AL, Foster AB, Del Rosso L, Mendoza SP, Capitano JP. Cortisol in mother's milk reflects maternal life history and predicts infant temperament. *Behav Ecol.* 2015;26:269-281.
- [118] Adcock IM, Ito K & Barnes PJ. Glucocorticoids; effects on gene transcription. *Proc Am Thor Soc.* 2004;1:247-254.
- [119] Weaver IC. Epigenetic effects of glucocorticoids. *Semin Fetal Neonatal Med.* 2009;14:143-150.
- [120] Wyrwoll CS, Mark PJ, Waddell BJ. Developmental programming of renal glucocorticoid sensitivity and the renin-angiotensin system. *Hypertension* 2007;50:579-584.
- [121] Wyrwoll CS, Kerrigan D, Holmes, MC, Seckl JR, Drake AJ. Altered placental methyl donor transport in the dexamethasone programmed rat. *Placenta* 2012;33:220-223.
- [122] Vaughan OR, Phillips HM, Everden AJ, Sferruzzi-Perri AN, Fowden AL. Dexamethasone treatment of pregnant F0 mice leads to parent of origin-specific changes in placental function of the F2 generation. *Reprod Fert Develop.* 2015;27:704-711.
- [123] Drake AJ, Liu L, Kerrigan D, Meehan RR & Seckl JR. Multigenerational programming in the glucocorticoid programmed rat is associated with generation-specific and parent of origin effects. *Epigenetics* 2011;6:1334-1343.
- [124] Long NM, Ford SP, Nathanielsz PW. Multigenerational effects of fetal dexamethasone exposure on the hypothalamic-pituitary-adrenal axis of the first and second generations female offspring. *Am J Obstet Gynecol.* 2012;208: 217 e1-8.
- [125] Long NM, Smith DT, Ford SP, Nathanielsz PW. Elevated glucocorticoids during ovine pregnancy increases appetite and produces glucose dysregulation and adiposity in their granddaughters in response to ad libitum feeding at 1 year of age. *Am J Obstet Gynecol.* 2013;209:353, e1-9.

Acknowledgements

We are grateful to all the members of the Department of Physiology, Development and Neuroscience who have helped with studies described here. We would also like to thank the BBSRC, the Horserace Betting Levy Board and the Centre for Trophoblast for their financial support.

Figure legends


Figure 1: Schematic diagram showing the sources of cortisol in the fetal circulation and the role of 11 beta-hydroxysteroid dehydrogenase as a placental barrier to materno-fetal cortisol transfer in sheep. Open arrows = major cortisol movements. Dashed arrow = minor cortisol movement.

→ Stimulatory effect. —| Inhibitory effect.

Data from references 14-18.


Figure 2: Fetal cortisol concentrations with respect to proximity to delivery in different species. Length of pregnancy: Pig 115 days (filled circles), Sheep 145 days (open circles), Human 280 days (filled triangle), Cow 280 days (filled squares), Horses (Pony) 335 days (open triangles). Data from references 1,22,23.


Figure 3: The endocrine systems affected by natural and synthetic glucocorticoids in fetal sheep together with the cellular and molecular processes within these endocrine systems influenced by prenatal glucocorticoid exposure. Data from references 1-11,42-44.



801
802
803
804
805
806
807
808
809
810
811
812
813

Figure 2





827

828

829

Table 1: The postnatal outcome of early life overexposure to glucocorticoids in farm animals.

	Species	Agent	Stage of pregnancy at overexposure	Postnatal outcomes	Age at outcome N = Neonate J = Juvenile A = Adult	Sex of Offspring M = Male F = Female	Reference
Maternal Synthetic Glucocorticoid Overexposure	Sheep	Dex	20%	↓ Birth weight and abdominal circumference	N	M	55
				Altered baroreflexes, ↑ sympathetic activity	N	M & F	56
				Altered vasodilatory responses	N	M & F	56
				Hypertension	J (4 mo)	M & F	57
					A (1.5-2 yr)	M & F	58
					A (5-6 yr)	F	59
					A (7 yr)	M & F	60
				Left ventricular hypertrophy ↑ Cardiac output	A (7 yr)	F	58
					A (7 yr)	F	58
				Altered endothelial superoxide production	J (4 mo)	M & F	61
				Altered cardiac mitochondrial function	J (6 mo)	M	62
				Altered brain RAS function ↑ brainstem AT1 receptor abundance	A (4-5 yr)	M	63
					A (7 yr)	F	60
				↓ Nephron number	A (5-6 yr)	F	59
					A (7 yr)	M & F	58
				↑ Mean glomerular volume	A (7 yr)	M & F	60
				↑ single nephron glomerular filtration rate	A (5-6 yr)	F	59
				↑ Glucose stimulated insulin secretion	A (4 yr)	M	54
				↑ Glucose tolerance	A (4 yr)	M	54
				↑ sensitivity to inhibition of lipolysis by insulin	A (5 yr)	F	64
			30%	↑ pituitary-adrenal responsive to stress	N (30 d)	F not M	65
				↓ liver, adrenal, pituitary & kidney weight	J (7 mo)	F	66
				↓ pituitary-adrenal responsiveness	J (7 mo)	F not M	67
				↑ Hippocampal GR mRNA	J (7 mo)	M not F	67
				↑ Hypothalamic AVP & GR mRNA	J (7 mo)	M not F	67
				↓ Pituitary POMC mRNA	J (7 mo)	F not M	67

				↑Adrenal ACTH receptor, StAR and 3βHSD mRNA	J (7 mo)	M not F	67
			70%	↓body weight and CRL	N	F	68
				↓basal and stimulated HPA function	A (2.5-3.5 yr)	F	69
				↓Glucose tolerance	A (2.5-3.5 yr)	F	69
				↓Insulin secretion	A (2.5-3.5 yr)	F	69
			70-82%	↓ brain weight and size	N	M & F	54
			98%	↑UCP1 content in brown adipose tissue	N	M & F	70
				↑Prolactin receptor abundance	N	M & F	70
				↑Relative fat mass	A (16 mo)	F not M	71
		Beta	55%	↑sympathetic and HPA responses	J (40 d)	F	72
				↑basal and ACTH stimulated cortisol secretion	A (1.5 yr)	F not M	73
				Hypertension	J (6 mo)	M & F	74
					A (1-2yr)	M	75
				Altered systemic and renal RAS function	J (6 mo)	M	76
				↓plasma renin and All concentrations	J (6 mo)	M & F	77
				↑All stimulated ROS production	J (6 mo)	M & F	76
				Altered renal All responsiveness	J (6 mo)	M & F	78
					A (1-1.5 yr)	M & F	79
				↓nephron number	J (6 mo)	M & F	80
				↓glomerular filtration rate	J (6 mo)	M	80
				Altered cerebral vascular tone and reactivity	A (1.5 yr)	F	81
				↑plasma leptin	A (1.5 yr)	M & F	73
				↑ leptin inhibition of adrenal function	A (1.5 yr)	M not F	73
			72-84%	↓Brain weight	J (6 wk)	M & F	66
					A (3.5 yr)	M & F	82
				↓Lung weight	J (12 wk)	M & F	66
				↓testicular development	J (6 & 12 wk)	M	83
				↓Body weight	J (12 wk)	M & F	84
				Hypotension	J (12 wk)	M & F	82
				↓plasma T ₃ levels	J (6 & 12 wk)	M & F	66
				↓plasma IGF-I & IGFBP levels	J (12 wk)	M & F	85

				↓hypothalamic AVP & CRH mRNA	J (6 & 12 wk)	M & F	65
				↓pituitary POMC, PC1 & PC2 mRNA	J (6 & 12 wk)	M & F	65
				↓ pituitary CRH/AVP responsiveness	J (7 mo)	F not M	67
				↓Pituitary GR	J (7 mo)	M & F	67
				↑basal and ACTH stimulated cortisol levels	A (1 yr)	M & F	86
				↑pituitary CRH/AVP responsiveness	A (2 yr)	M & F	86
				↓adreno-cortical ACTH responsiveness	A (3 yr)	M & F	87
				↓basal ACTH and cortisol levels	A (3 yr)	M & F	87
				↓plasma glucose levels	J (12 wk)	M & F	65
				Insulin resistance	J (6 mo)	M & F	84
				Glucose intolerant	A (1.5 yr)	M & F	84
				↑Glucose stimulated insulin secretion	A (1.5 yr)	M & F	84
				↑Fasting insulin:glucose ratio	A (2 & 3 yr)	M & F	88
				↑Hepatic glucose-6-phosphatase activity	A (3.5 yr)	M & F	88
	Horse	Dex	95%	↓Body weight	N	M & F	89
				↓ CRL and adreno-cortical ACTH responsiveness	N	M & F	90
Maternal Cortisol Overexposure	Sheep	Cortisol	20%	↑renal Na ⁺ -K ⁺ ATPase α-subunit	J (2 mo)	M & F	59
				↓glomerular number	A (4-5 yr)	F	59
				↑single nephron GFR	A (4-5 yr)	F	59
				Hypertension	A (1.5 yr)	M & F	63
					A (4-5 yr)	F	59
				Fasting hyperinsulinaemia	A (4 yr)	M	54
				↑Glucose stimulated insulin secretion	A (4 yr)	M	54
			50-72%	↓Body weight	N	M & F	90
		Periodic isolation	72%-term	↑Birth weight	N	M & F	92
				↑Basal Cortisol concentration	N	M & F	92
	Pig	Cortisol	33-50%	↑body weight	J (5 mo)	M	93
		ACTH	40%	↑basal LH	N	F	94
			40-65%	↑plasma CBG & ↑adrenomedullary cells	N	M & F	95
			40-73%	↑adrenal cortex:medulla ratio	N & J (60 d)	M & F	96
				↑Hypothalamic CRH and adrenal ACTH receptor	N	M & F	96
				↓Hypothalamic endorphin	J (30 d)	M & F	96

				↑Pituitary POMC mRNA	J (60 d)	M & F	96
				↑HPA stress responsiveness	J (11 wk)	F	96
			75-93%	↓Body weight	N & J	M & F	95
				↑plasma CBG & 5HT	N & J	M & F	95
				↓Relative adrenal weight ↑adrenal cortex area	J (4 wk)	M & F	95
				Altered brain neurotransmitter system	J (4 wk)	M & F	95
		Social mixing	35-56%	↑hypothalamic CRH expression to social stress	J (9 wk)	F	97
				↑Cortisol response to social stress	J (9 wk)	F	97
			67-97%	↑Cortisol response to social stress	J (9 wk)	F	97
	Cow	ACTH	20-50%	↑Body weight	N (at birth)	M & F	98
				↑Cortisol secretion to restraint	J (5 mo)	M & F	98
	Transport	20-50%		↑Cortisol secretion to restraint	J (5 mo)	M & F	98
				↓Cortisol clearance	J (5 mo)	M & F	98
				↑Heart rate increment to restraint	J (5 mo)	M & F	98
Fetal Synthetic Glucocorticoid Overexposure	Sheep	Beta	72-84%	↑Glucose-stimulated insulin secretion	J (6 mo)	M & F	83
				Glucose intolerant	A (1 yr)	M & F	83
				↓Basal insulin concentration	A (2 yr)	M & F	88
				↑Basal insulin concentration	A (3 yr)	M & F	88
				↑Hepatic glucose-6-phosphatase activity	A (3.5 yr)	M & F	88
				↓ Pituitary CRH/AVP responsiveness	A (1 yr)	M & F	86
				↑ Adreno-cortical ACTH responsiveness	A (1 yr)	M & F	86
				↓Basal and stimulated ACTH concentration	A (2 yr)	M & F	87
				↑ Adreno-cortical ACTH responsiveness	A (2 yr)	M & F	87
				↓Brain weight	A (3.5 yr)	M & F	82
Neonatal Glucocorticoid Overexposure	Sheep	Dex	3-4 days	Altered NMDA receptor kinetics	5 d	M & F	99
	Horse	ACTH	1-5 days	↓Glucose stimulated insulin secretion	J (2 & 12 wk)	M & F	100
				↑Basal cortisol concentrations	J (12 wk)	M & F	101
				Altered pituitary sensitivity to hypoglycaemia	A (1 & 2 yr)	M & F	102

830

831 Dex=dexamethasone, Beta=betamethasone, ACTH=Adrenocorticotrophic hormone, All=Angiotensin II, AT1=Angiotensin receptor type 1, AVP=Arginine
832 vasopressin, CBG=Corticosteroid binding globulin, CRH=Corticotropin releasing hormone, CRL=Crown rump length, GR= Glucocorticoid receptor,
833 GRF=glomerular filtration rate, HPA=hypothalamic-pituitary-adrenal, 3 β HSD=hydroxysteroid dehydrogenase, 5HT=5-hydroxytryptamine, IGF-I=Insulin-like
834 growth factor I, IGFBP=Insulin-like growth factor binding protein, LH=Luteinising hormone, NMDA=N-methyl-D-aspartate, POMC=Pro-opiomelanocortin,
835 RAS=Renin-angiotensin system, ROS=Reactive oxygen species, StAR=Steroidogenic acute regulatory protein, T₃=Tri-iodothyronine, UCP1=Uncoupling
836 protein 1.

837

838

839

840

841

842

843

844

845

846

847

848

